


INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification : C12N 15/00, C12P 21/04 C07K 13/00	A1	(11) International Publication Number: WO 90/08187 (43) International Publication Date: 26 July 1990 (26.07.90)
(21) International Application Number: PCT/US89/00218 (22) International Filing Date: 19 January 1989 (19.01.89) (71) Applicant (for all designated States except US): DANA FARBER CANCER INSTITUTE [US/US]; 44 Binney Street, Boston, MA 02115 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): REINHERZ, Ellis, L. [US/US]; 113 South Great Road, Lincoln, MA 01773 (US). SAYRE, Peter, H. [US/US]; 64 Richardson Street, New Brunswick, NJ 08901 (US). HUSSEY, Rebecca, E. [US/US]; 113 South Great Road, Lincoln, MA 01773 (US). CHANG, Hsiu-Ching [— /US]; 86 Emerson Gardens Road, Lexington, MA 02173 (US).	(74) Agents: DECONTI, Giulio, A., Jr. et al.: Hamillon, Brook, Smith & Reynolds, Two Militia Drive, Lexington, MA 02173 (US). (81) Designated States: AT (European patent), BE (European patent), CH (European patent), DE (European patent), FR (European patent), GB (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent), US. Published <i>With international search report.</i>  Document provided by: Ralph McElroy Translation Company Call 1-800-531-9977 (outside Texas) or (512) 472-6753 (inside Texas)	
(54) Title: SOLUBLE TWO DOMAIN CD2 PROTEIN (57) Abstract A soluble CD2 peptide having an LFA-3 binding domain and antigenic epitopes recognized by antibodies raised against native CD2 on the surface of T cells and is capable of forming at least two intramolecular disulfide bonds. The peptide is capable of binding LFA-3, inhibiting CD2-mediated T cell activation and reacting with antibodies raised against CD2. Preferably, the soluble peptide comprises an amino acid sequence of 182 residues in length which is encoded by two extracellular segment exons of the gene encoding human CD2 glycoprotein.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MR	Mauritania
BE	Belgium	GA	Gabon	MW	Malawi
BF	Burkina Faso	GB	United Kingdom	NL	Netherlands
BG	Bulgaria	HU	Hungary	NO	Norway
BJ	Benin	IT	Italy	RO	Romania
BR	Brazil	JP	Japan	SD	Sudan
CA	Canada	KP	Democratic People's Republic of Korea	SE	Sweden
CF	Central African Republic	KR	Republic of Korea	SN	Senegal
CG	Congo	LI	Liechtenstein	SU	Soviet Union
CH	Switzerland	LK	Sri Lanka	TD	Chad
CM	Cameroon	LU	Luxembourg	TG	Togo
DE	Germany, Federal Republic of	MC	Monaco	US	United States of America
DK	Denmark				

-1-

SOLUBLE TWO DOMAIN CD2 PROTEINBackground

The human CD2 (T11) molecule is a 50KD surface glycoprotein expressed on >95% of thymocytes and virtually all peripheral T lymphocytes which mediates both adhesion between these cells and their cognate partners as well as subsequent activation events. Specific combinations of antibodies against the surface-bound molecule can activate IL-2 dependent T cell proliferation, helper T cell function and cytotoxicity by natural killer cells and cytolytic T lymphocytes (Meuer, S.C., et al., Cell 36:897-906 (1984); Brottier, P., et al., J. Immunol., 135:1624-1631 (1985); Siliciano, R.F., et al., Nature 317:428-430 (1985)) in the absence of cellular adhesion. In addition, thymocyte activation can be mediated via CD2 (Fox, D.A., et al., J. Immunol. 134:330-335 (1985); Denning, S.M., et al., J. Immunol. 139:2573-2578 (1987)). The role of CD2 in approximation of T cells to various cell types including human thymic epithelial cells, B cells, target cells and sheep erythrocytes has been demonstrated to depend on direct interaction between CD2 and the broadly distributed human lymphocyte function-associated antigen 3 (LFA-3) surface glycoprotein or its sheep homologue, T11TS (Denning, S.M., et al., J. Immunol. 139:2573-2578 (1987); Hunig, T., J. Exp. Med. 162:890-901 (1985); Hunig, T., et al., Nature 326:298-301 (1987); Shaw, S. et al., Nature 323:262-264 (1986); Selvaraj, P., et al., Nature 326:400-403 (1987); Vollger, et al., J. Immunol. 138:358-363 (1987)).

Biochemical analyses using specific monoclonal antibodies show that CD2 is T lineage-specific and exists on the cell surface in several differentially glycosylated forms (Howard, F.D., et al., J. Immunol. 126:2117-2122 (1981); Brown, M.H., et al., In Leucocyte Typing III ed McMichael, A.J., Oxford University Press pp. 110-112 (1987); Sayre, P.H., et al., Proc. Natl. Acad. Sci. USA 84:2941-2945 (1987)). CD2 cDNA clones predict a cleaved signal peptide of 24 amino acid residues, an extracellular segment of 185 residues, a transmembrane domain of 25 residues and a cytoplasmic region of 117 residues (Sayre, P.H., et al., Proc. Natl. Acad. Sci. USA 84:2941-2945 (1987); Sewell, W.A., et al., Proc. Natl. Acad. Sci. USA 83:8718-8722 (1986); Seed, B. and A. Aruffo, Proc. Natl. Acad. Sci. USA 84:3365-3369 (1987); Clayton, L.K., et al., Eur. J. Immunol. 17:1367-1370 (1987)). The corresponding genomic organization reveals a single exon encoding the signal peptide (less four residues), two exons encoding the extracellular segment, one exon encoding the transmembrane domain and charged membrane anchor segment, and one exon encoding the cytoplasmic region (Diamond, D.J., et al., Proc. Natl. Acad. Sci. USA 85:1615-1619 (1988)).

Summary of the Invention

This invention pertains to a soluble peptide having a lymphocyte function-associated antigen 3 (LFA-3) binding domain and antigenic epitopes recognized by antibodies raised against native

surface CD2 on resting T lymphocytes. The soluble peptide is capable of forming at least two intramolecular disulfide bonds and binds the surface-bound CD2 legand, LFA-3. The soluble peptides of this invention exist as monomers in an aqueous medium. Preferably, the soluble peptides have an amino acid sequence encoded by the two extracellular segment exons of the gene encoding human CD2 glycoprotein. Most preferably, the soluble peptides have about 182 amino acid residues encoded by the two extracellular segment exons of the gene encoding human CD2 glycoprotein. Soluble peptides of this invention can be made by enzymatic fragmentation, peptide synthesis, or recombinant DNA technology. They can be used to block T cell function which is dependent upon antigen activation.

Brief Description of the Drawings

Figure 1 shows the amino acid sequence and exon organization of human CD2.

Figure 1a shows the DNA sequence of human CD2 glycoprotein.

Figure 2 shows the structure of recombinant soluble T11_{ex2}, native CD2 and its genomic organization.

Figure 3 shows the construction of expression plasmid for production of the CD2 external segment molecule T11_{ex2}.

Figure 4 shows SDS-PAGE analysis of purified, radioiodinated and endoglycosidase digested T11_{ex2}.

Figure 5 shows equilibrium sedimentation data.

Figure 6 shows the circular dichroism spectra of T11_{ex2}.

Figure 7 shows the competitive inhibition of radioiodinated T11_{ex2} binding of JY cells.

Figure 8 shows the saturation binding of T11_{ex2} to JY cells and Scatchard analysis.

Detailed Description of the Invention

This invention pertains to a soluble peptide having a lymphocyte function-associated antigen 3 (LFA-3) binding domain and antigenic epitopes recognized by antibodies raised against native surface CD2 on resting T lymphocytes. The soluble peptide is capable of forming at least two intramolecular disulfide bonds and binds the surface-bound CD2 ligand, LFA-3. Specifically, the antigenic epitopes which are localized on the peptide are T11₁, T11₂ and T11₃. The soluble peptides are further capable of inhibiting CD2-mediated T cell activation and are soluble in aqueous medium and exist as monomers in the same.

In one embodiment, the soluble peptides have an amino acid sequence encoded by the two extracellular segment exons of the gene encoding human CD2 glycoprotein. The soluble peptides are capable of binding the surface-bound CD2 ligand, LFA-3, and can react with antibodies raised against native CD2 on the surface of human T cells. The soluble peptides are further capable of inhibiting CD2-mediated T cell activation. The peptides exist as monomers in an aqueous medium.

In the preferred embodiment, the soluble peptide comprises about 182 amino acid residues

-5-

encoded by the two extracellular segment exons of the gene encoding human CD2 glycoprotein. The soluble peptide is monomeric in aqueous medium and comprises four cysteine residues located in the carboxy-terminal region of the amino acid sequence. The peptide is capable of forming at least two sets of intramolecular disulfide bonds between the amino-terminal cysteines and the carboxy-terminal cysteines. Preferably, the peptide has the following amino acid sequence:

amino

```

      Lys Glu Ile Thr Asn Ala Leu Glu Thr Trp
11  Gly Ala Leu Gly Gln Asp Ile Asn Leu Asp
21  Ile Pro Ser Phe Gln Met Ser Asp Asp Ile
31  Asp Asp Ile Lys Trp Glu Lys Thr Ser Asp
41  Lys Lys Lys Ile Ala Gln Phe Arg Lys Glu
51  Lys Glu Thr Phe Lys Glu Lys Asp Thr Tyr
61  Lys Leu Phe Lys Asn Gly Thr Leu Lys Ile
71  Lys His Leu Lys Thr Asp Asp Gln Asp Ile
81  Tyr Lys Val Ser Ile Tyr Asp Thr Lys Gly
91  Lys Asn Val Leu Glu Lys Ile Phe Asp Leu
101 Lys Ile Gln Glu Arg Val Ser Lys Pro Lys
111 Ile Ser Trp Thr Cys Ile Asn Thr Thr Leu
121 Thr Cys Glu Val Met Asn Gly Thr Asp Pro
131 Glu Leu Asn Leu Tyr Gln Asp Gly Lys His
141 Leu Lys Leu Ser Gln Arg Val Ile Thr His
151 Lys Trp Thr Thr Ser Leu Ser Ala Lys Phe
161 Lys Cys Thr Ala Gly Asn Lys Val Ser Lys
171 Glu Ser Ser Val Glu Pro Val Ser Cys Pro
181 Glu Lys

```

carboxy

-6-

The soluble peptide of about 182 amino acid residues in length has an LFA-3 binding domain and comprises the antigenic epitopes T11₁, T11₂ and T11₃ which are localized on the soluble peptide. The epitopes, which are localized on this peptide are antigenic determinants which are recognized by antibodies against the native surface CD2 structure on resting T lymphocytes. Thus, the peptides of this invention can be used to inhibit CD2-mediated T cell activation.

The above amino acid sequence corresponds with a portion of the naturally occurring extracellular domain of CD2 that is capable of binding LFA-3, inhibiting CD2-mediated T cell activation and reacting with antibodies raised against native CD2 on the surface of human T cells. Amino acid sequences embraced by this invention include analogous or homologous sequences which encode proteins capable of binding LFA-3, inhibiting CD2-mediated T cell activation and reacting with antibodies raised against native CD2. In addition, the peptide structure can be modified by deletions, additions, inversions, insertions or substitutions of one or more amino acid residues in the sequence to yield peptides having the characteristics of the peptides of this invention. All such modifications of the above amino acid sequence are embraced by this invention without essentially detracting from the properties of the peptide, i.e., the capacity of the peptide to bind LFA-3, inhibit CD2-mediated T cell activation and react with antibodies raised against native CD2 on the surface of human T cells.

-7-

Naturally occurring allelic variations and modifications are included within the scope of the invention so long as the variation does not substantially reduce the ability of the peptide to bind LFA-3 and inhibit CD2-mediated T cell activation.

The soluble peptides of this invention carry epitopes recognized by antibodies against native surface CD2 structure on resting T lymphocytes and interacts specifically with the surface-bound CD2 ligand, LFA-3. Additionally, the soluble peptide exists as a monomer in aqueous medium and includes a proteolytically-resistant amino-terminal fragment encoded by the first extracellular segment exon of the gene encoding human CD2 glycoprotein. The proteolytically resistant fragment comprises about 100 amino acid residues which correspond with the amino-terminal portion of the extracellular domain of human CD2 glycoprotein and is capable of inhibiting CD2-mediated T cell activation. The 100 amino acid fragment has been described in U.S. Patent Application Serial No. 07/293,330 filed January 4, 1989, by Reinherz et al., the teachings of which are incorporated herein by reference. The entire amino acid sequence and exon organization of human CD2 glycoprotein is shown in Figure 1. This amino acid sequence of human CD2 was deduced from cDNA (Sayre, P.H., et al., Proc. Natl. Acad. Sci. USA 84:2941-2945 (1987); Sewell, W.A., et al., Proc. Natl. Acad. Sci. USA 83:8718-8722 (1986); Seed, B. and A. Aruffo, Proc. Natl. Acad. Sci. USA 84:3365-3369 (1987)) and genomic clones (Diamond, et al., Proc. Natl. Acad. Sci. USA 85:1615-1619 (1988)).

-8-

This invention also pertains to an isolated DNA sequence that encodes a soluble monomeric human CD2 peptide having an LFA-3 binding domain and antigenic epitopes recognized by antibodies raised against native CD2 on the surface of human T cells and is capable of forming at least two intramolecular disulfide bonds. Specifically, the antigenic epitopes are T11₁, T11₂ and T11₃. The peptide that is encoded by the isolated DNA of this invention and is capable of binding LFA-3, inhibiting CD2-mediated T cell activation and reacting with antibodies raised against native CD2 on the surface of human T cells. In a preferred embodiment, the isolated DNA sequence encodes a soluble peptide having an amino acid sequence encoded by the two extracellular segment exons of the gene encoding the human CD2 glycoprotein. The encoded peptide is capable of binding LFA-3, inhibiting CD2-mediated T cell activation and reacting with antibodies raised against native CD2. Preferably, the isolated DNA of the invention encodes the 182 amino acid sequence shown above or substantial coding equivalent thereof. The DNA sequence can be modified by deletion, insertion or substitution of nucleotides to yield peptides which exhibit substantially the same properties of the above peptide of about 182 amino acid residues. All such modifications of the DNA sequence are within the scope of this invention so long as the DNA sequence encodes a soluble peptide that is capable of binding LFA-3 and inhibiting CD2-mediated T cell activation. The isolated DNA sequences of the invention can be made

using recombinant DNA technology or chemically synthesized. The DNA sequence for CD2 glycoprotein is shown in Figure 1a.

This invention further pertains to a recombinant expression vector comprising the DNA sequence encoding a soluble, monomeric human CD2 peptide having an LFA-3 binding domain and the antigenic epitopes T11₁, T11₂ and T11₃. The antigenic epitopes are recognized by antibodies raised against native CD2 on resting T cells and interacts specifically with LFA-3. The encoded soluble protein is capable of forming at least two intramolecular disulfide bonds. Alternatively, the expression vector comprises a DNA sequence encoding a soluble CD2 protein encoded by the two extracellular segment exons of the gene encoding human CD2 which is capable of inhibiting CD2-mediated T cell activation and LFA-3 binding. Preferably, the expression vector is a baculovirus transfer vector and comprises a DNA sequence which encodes a peptide of about 182 amino acid residues as shown above. Other vectors, however, may be used, including prokaryotic and eukaryotic expression systems. Modifications of the peptide in which amino acid residues have been deleted, inserted or substituted without essentially detracting from the properties of the peptide are embraced by the invention. The invention further pertains to cells transformed with the above expression vector.

Soluble peptides of this invention can be made by enzymatic fragmentation of human CD2 glycoprotein

or a portion thereof, by peptide synthesis or recombinant DNA technology. Preferably, the soluble CD2 peptides will be produced by inserting DNA encoding a peptide sequence which is capable of binding LFA-3 and inhibiting CD2-mediated T cell activation (e.g., CD2 DNA which represents the desired amino acid sequence of the extracellular domain of CD2) into an expression vector. The transformed cells then express the soluble human CD2 peptide encoded by the two extracellular segment exons of the gene encoding the human CD2 glycoprotein. The peptide expressed by the transformed cells is capable of binding LFA-3 and inhibiting CD2-mediated T cell activation. In addition to genetic engineering techniques for synthesizing soluble peptides of the invention, the soluble peptides can be synthesized directly by procedures of chemical protein synthesis. For example, the above 182 amino acid sequence or modified equivalent thereof can be synthesized by the solid phase procedure of Merrifield.

This invention further pertains to a method of inhibiting T cell activation, comprising the step of administering to a patient, a soluble peptide having an LFA-3 binding domain, antigenic epitopes recognized by antibodies raised against native CD2 on the surface of human T cells and is capable of forming at least two intramolecular disulfide bonds. In another embodiment, a patient is administered a solution containing a soluble peptide having an amino acid sequence encoded by the two extracellular

-11-

segment exons of the gene encoding human CD2 glycoprotein. The peptide is capable of binding LFA-3, inhibiting T cell activation and reacting with antibodies raised against the native CD2 protein. Preferably, the soluble peptide comprises a sequence of about 182 amino acid residues which corresponds with the portion of the extracellular domain of human CD2 glycoprotein or fragment thereof which is capable of binding LFA-3 and inhibiting CD2-mediated T cell activation. In one embodiment of the method, the soluble peptide can be administered intravenously.

The soluble CD2 peptides of this invention generally will bind to human lymphocytes and human red blood cells which express a homologous set of surface structures. The soluble peptides are also capable of competing with the naturally-present CD2 on the surface of a human lymphocyte, thus interfering with the ability of the lymphocyte to make contact with its target cells (if the lymphocyte is a cytolytic cell), or with macrophages having CD2 binding structures which permit the cell-to-cell contact necessary for lymphocyte proliferation. To test a soluble peptide for the ability to inhibit lymphocyte proliferation, or the cytotoxic effector function, the soluble peptide is contacted with the lymphocytes prior to stimulation with mitogen, and degree of proliferation is measured, using standard techniques, and the result compared to a control in which the soluble peptide was not used.

-12-

The soluble peptides of this invention can be used in a variety of diagnostic and therapeutic applications in which the CD2 surface glycoprotein is expressed on the surface of many human T cell malignancies, e.g., T cell leukemias and lymphomas. In addition, autoimmune diseases, e.g., rheumatoid arthritis and Systemic Lupus Erythematosus (SLE), are characterized by the presence in the blood and lymph of large numbers of CD2-bearing T cells. Rapid cell turnover in these disease states can cause the shedding of the CD2 molecule into the bloodstream.

The CD2 soluble peptides of this invention can be used as an immunogen to produce polyclonal or monoclonal anti-CD2 antibodies, using conventional techniques. These antibodies can be labeled with any conventional label, e.g., radioisotopes, and used in conventional immunoassay methods to measure serum CD2 levels and thus monitor patients having T cell associated diseases. Particularly sensitive ELISA-type assays will employ two anti-CD2 antibodies, each to a different antigenic determinant on the surface of CD2, in a sandwich format.

The disease states which can be treated using the soluble peptides of this invention include medical conditions characterized by unwanted activity of the immune system which results in excess T cell activation, which plays a key role in the amplification of the immune response. These conditions include SLE; juvenile onset diabetes; multiple sclerosis; allergic conditions; inflammatory conditions such as exzema, ulcerative

-13-

colitis, inflammatory bowel disease, and Crohn's disease; and allograft rejection (e.g., rejection of a transplanted heart or kidney). The soluble CD2 peptides compete with the surface-bound CD2 for its ligand on target cells thus dampening immune response amplification. The soluble CD2 peptide admixed with a pharmaceutically acceptable carrier substance such as saline, is administered intravenously to a human patient in an effective amount, e.g., 20 μ g to 500 μ g per kg body weight. Preferably, the soluble peptide admixed with a pharmaceutically acceptable carrier comprises about 182 amino acid residues which correspond to a portion of the extracellular domain of CD2 that is capable of binding LFA-3 and inhibiting CD2-mediated CD2 activation. For some conditions, a soluble CD2 peptide can be administered directly to the site where needed most; for example, a soluble CD2 peptide can be injected directly into the inflamed joint of a human patient suffering from rheumatoid arthritis.

This invention is further illustrated by the following Exemplification.

EXEMPLIFICATION

The term T11_{ex2}, as referred to in the Exemplification, is defined herein to be a soluble protein having an amino acid sequence encoding the two extracellular segment exons and a codon (182-Lys) derived from the transmembrane domain exon of the CD2 glycoprotein. The entire amino acid

-14-

sequence for CD2 has been described. See U.S. Patent Application Serial No. 932,871, filed November 18, 1986, by Reinherz et al., the teachings of which are incorporated herein by reference.

Baculovirus Transfer Vector Plasmid Construct for Tl1_{ex2}

The plasmid pAc373/Tl1_{ex2} was constructed by digestion of pGEM-4-S1, a pGEM derivative containing a 950 bp fragment of the CD2 cDNA PB2 (Sayre et al., Proc. Natl. Acad. Sci. USA 84:2941-2945 (1987)) with PvuII, which digests the cDNA at nucleotide position 628 near the start of the transmembrane region. A double-stranded synthetic oligonucleotide linker:

CTGTCCAGAGAAATAAGGATCCT
GACAGGTCTCTTTATTCTAGGA

containing the last base for the Ser-178 codon, codons for Cys-Pro-Glu-Lys, a stop codon and a BamHI restriction site was synthesized and ligated to the blunt PvuII ends. After BamHI digestion, the insert was cloned into the BamHI site of the pAc373 baculovirus transfer vector (Smith et al., Annu. Rev. Immuno. 2:319-333 (1985)). All restriction enzymes were produced from New England Biolabs (Beverly, MA).

Recombinant protein production

Transfer of the Tl1_{ex2} sequences from the plasmid vector to the AcNPV genome to generate recombinant baculovirus Tl1_{ex2}-AcNPV was ac-

complished essentially as described (Smith et al., Proc. Natl. Acad. Sci. USA, 82:8404-8408 (1985); Hussey et al., Nature, 331:78-81 (1988). Metabolic labeling with ^{35}S -cysteine of T11_{ex2}-AcNPV-infected SF9 cells was carried out as described (Hussey et al., Nature 331:78-81 (1988). Metabolically labeled culture supernatants were harvested, microfuged for 10 min. and subjected to immunoprecipitation for 16 h at 4°C with a monoclonal anti-CD2 antibody (anti-T11₁, 3T4-8B5) linked to Affigel-10 (Biorad) beads (10 mg monoclonal antibody/ml gel). After immunoabsorption the beads were washed twice with lysis buffer and bound material eluted with 0.1 M glycine, HCl buffer, pH2. Eluates were analyzed by SDS-PAGE in 12.5% gels. Large protein production was performed as described (Hussey et al., Nature 331:78-81 (1988)) except that proteins were purified over an anti-T11₁ (3T4-8B5) immunoaffinity column.

Protein microsequencing

Proteins were prepared for microsequencing by electrophoresis on 12.5% polyacrylamide gels, followed by electroblotting onto polyvinylidene difluoride membranes according to the method of Matsudaira, J. Biol. Chem. 262:10035-10038 (1987). After visualization with Coomassie blue, stained bands were excised and loaded onto an Applied Biosystems (Foster City, CA) 470A sequencer and sequenced using the O3RPTH program.

Endoglycosidase F digestion of T11_{ex2}

1 μ g samples of purified T11_{ex2} dialyzed against PBS were incubated in Endo-F buffer (0.1 M sodium phosphate, 1% NP-40, 1% 2-mercaptoethanol, 50 mM EDTA, 1 mM phenylmethyl-sulfonyl fluoride, pH 6.1) in the presence of 0.7 units Endo-F (NEN) in 7 μ l reaction volumes. Digestion was arrested at the indicated times by addition of 15 μ l SDS sample buffer, boiling for 5 min. and freezing at -20°C prior to analysis by 12.5% SDS-PAGE.

Equilibrium sedimentation

Sedimentation studies were performed using the short column, high speed meniscus-depletion method of Yphantis Biochem. 3:297-317 (1964); and Richardson et al., Biochem. J. 135:87-92 (1973)). Standard double-sector cells, equipped with sapphire windows, 4 mm solution column lengths and a temperature of 21°C were used.

Papain digestion

Papain (32 ng) was added to 8 μ g samples of T11_{ex2} at 0.5 mg/ml in PBS containing 10 mM DTT for an enzyme:protein ratio of 1:250. Samples were incubated at 37°C for 15, 30, 45 or 60 min. Digestions were stopped by the addition of SDS buffer and boiling for 5 min. Samples were electrophoresed on a 12.5% polyacrylamide gel and stained with Coomassie blue.

Circular dichroism

Far ultraviolet CD spectra were obtained on an Instruments SA Jobin Yvon circular dichrograph calibrated with (+)10-camphorsulfonic acid and epiandrosterone. Measurements were taken at 25, 50 and 80°C \pm 0.1°C in 10 mM sodium phosphate pH 7.2 in a 1 mm cell. All spectra represent an average of 3 to 5 individual spectra with data taken at 0.5 nm intervals using a 10 second response time for each point. Protein concentrations were determined by quantitative amino acid analysis of aliquots taken from the sample cells.

Rosette inhibition assay

Jurkat cells were washed with SMEM/2% FCS (wash) and resuspended at 10^7 /ml. Sheep erythrocytes were washed twice in HBSS and resuspended in wash to 5% v/v. 10 μ l of sheep erythrocytes were aliquoted into 12-75 mm plastic tubes and 100 μ l wash, T11_{ex2} protein or control soluble CD4 T4_{ex1} protein (Hussey et al., Nature 331:78-81 (1988)) was added, followed by incubation at 4°C for 30 min. Subsequently, 20 μ l of Jurkat cells were added after 5 min., centrifuged at 800 rpm in a Sorvall RT6000, followed by incubation at 4°C for 1 h. The cell mixture was gently resuspended and mounted on glass slides with cover slips and rosette formation assessed on a Zeiss photomicroscope.

Radiolabeled T11_{ex2} binding assays

The purified recombinant soluble CD2 extracellular domain molecule T11_{ex2} was radioiodinated as follows: 50 μ l T11_{ex2} (1 mg/ml) dialyzed against PBS was labeled with 10 μ l immobilized lactoperoxidase/ glucose oxidase (Enzymobeads; Biorad Laboratories, Richmond, CA) in 40 mM sodium phosphate, pH 7.2, 0.4% glucose and 1 mCi ¹²⁵I for 5 min. After quenching the reaction for 20 min. with 20 mM sodium iodide and 0.02% sodium azide, 20 μ l FCS was added and the free iodine separated on a 1 ml Bio-Gel P-6 column (Biorad Laboratories) conditioned with 0.2M sodium phosphate pH 7.2, 10% FCS and run in the same buffer.

Cold competition studies: 5 x 10⁶ cpm radio-labelled T11_{ex2} (2.8 x 10⁸ cpm/nmole) was added at 0.1 μ M to 1.8 x 10⁶ JY cells overlaid onto 0.2 ml of a 1.5:1 mixture of dibutyl phthalate:diethyl phthalate (Aldrich Chemical Co., Milwaukee, WI) in 0.5 ml plastic tubes as described in Teshigawara et al., J. Exp. Medicine 165:223-238 (1987).

Increasing concentrations of unlabelled T11_{ex2} or T4_{ex1} were added in a final volume of 200 μ l in RPMI 1640/10% FCS. After 1 h incubation at 4°C, the tubes were centrifuged (8,500 g for 1 min.), the tips of the tubes containing the cell pellets were cut, and the cell-bound and free radioactivity were determined in a gamma counter. To some tubes, anti-LFA-3 antibody TS2/9 (generously provided by Dr. Timothy Springer, Dana Farber Cancer Institute)

-19-

was added at 50 $\mu\text{g/ml}$ as a separate determination of nonspecific binding. This concentration of anti-LFA-3 was independently shown to give maximal inhibition of T11_{ex2} binding. Specific activity was calculated using a MW for T11_{ex2} of 30,000. Protein concentrations were determined by quantitative amino acid analysis of standard T11_{ex2} samples. Subsequent samples were compared to standards by Coomassie blue staining of two-fold dilutions of standard and test samples run on the same gel and analyzed by densitometry.

Saturation binding studies: Increasing concentrations of radiolabelled T11_{ex2} (1.31×10^7 cpm/nmole) were added to 2.6×10^6 JY cells in the presence or absence of 50 $\mu\text{g/ml}$ anti-LFA-3 antibody to determine nonspecific binding. Binding was carried out as above and the dissociation constant determined by Scatchard analysis after subtraction of nonspecific binding determined in the presence of anti-LFA-3.

RESULTS

Production and purification of T11_{ex2}

A construct for expression of a soluble fragment of CD2 that included all the residues encoded by the leader and two extracellular segment exons was designed (Figure 2, exons 1-3).

Figure 2 shows a comparison of the 182 extracellular CD2 amino acids comprising T11_{ex2} (top) to CD2 protein structure (middle). The positions of cysteine residues (C), carbohydrate addition sites

-20-

(CHO), the CD2 leader segment (L) and the CD2 transmembrane domain (TM) are indicated. In the CD2 gene (bottom), exon 1 corresponds to CD2 amino acid residues -24 to -4, exon 2 to residues -4 to 104, exon 3 to residues 104 to 181, exon 4 to residues 181 to 222, and exon 5 to residues 222 to 327 (Diamond *et al.*, PNAS USA 85:1615-1619 (1988)).

The plasmid pAc373/T11_{ex2} was constructed and encodes 182 amino acids of the predicted CD2 external segment including all of the residues derived from the two extracellular exons (Figure 2) and part of one codon (for Glu-181) and all of a second codon (for Lys-182) derived from the transmembrane domain exon. This construction, thus, includes all four extracellular cysteine residues located in domain II of CD2 and thereby avoids problems associated with intermolecular disulfide exchange observed with a previous construction (Richardson, *et al.*, Proc. Natl. Acad. Sci USA 85:5176-5180 (1988)).

Plasmid pAc373/T11_{ex2} was used to co-transfect SF9 with AcNPV baculoviral DNA. Recombinant baculovirus, termed T11_{ex2}-AcNPV, were selected, purified and used to infect small-scale cultures for metabolic labeling. Immunoprecipitation of radio-labelled supernatants with anti-T11₁ (3T4-8B5), an anti-CD2 specific monoclonal antibody (Meuer *et al.*, Cell 36:897-906 (1984)), verified that T11_{ex2}-AcNPV directed the production of a recombinant CD2 molecule in SF9 cells (data now shown).

T11_{ex2}-AcNPV was therefore used to infect liter cultures for the production of large amounts of protein. T11_{ex2} protein was purified from infected

-21-

cell supernatants by affinity chromatography on an anti-T11 column.

The construction of expression plasmid for production of the CD2 external segment molecule T11_{ex2} is shown in Figure 3. The plasmid pGEM-4-S1 carries a 950 bp fragment of the CD2 cDNA. After digestion of pGEM-4-S1 with PvuII, ligation of the double-stranded linker and further digestion with BamHI, the cDNA insert was isolated and ligated into the BamHI-digested baculoviral transfer vector pAc373. The resulting plasmid pAc373/T11_{ex2} encodes 182 amino acids of the mature CD2 extracellular segment. The promoter for the polyhedrin gene in the pAc373 transfer vector is shown by the black box and the polyhedrin gene is indicated by the open box. The 950 bp CD2 coding fragment in pGEM-4-S1 is shown in a stippled box. The position of the T7 polymerase promoter in the pGEM vector is shown.

Biochemical characterization of T11_{ex2}

Figure 4 shows the purification, radioiodination and endoglycosidase digestion of T11_{ex2}. Lanes a-d contain 1 μ g T11_{ex2} purified from large scale cultures of SF9 cells infected with T11_{ex2}-AcNPV was analyzed by Coomassie staining on a 12.5% polyacrylamide gel in the presence of 50 mM DTT (lane a) or in nonreducing conditions (lane b). An aliquot of T11_{ex2} radioiodinated with solid-phase lactoperoxidase/glucose oxidase was analyzed on the same gel in the presence (lane c) or absence (lane d) of 50 mM DTT by autoradiography. Lanes e-k

contain 1 μ g purified T11_{ex2} which was digested with 9.7 units Endo-F for varying amounts of time and the reaction stopped by the addition of SDS sample buffer prior to SDS-PAGE analysis and Coomassie staining. Lane e contains no enzyme; f, simultaneous addition of enzyme and sample buffer; g-j, 1 min, 5 min, 1 h, 18 h digestion; k, 0.7 units Endo-F alone.

T11_{ex2} migrates as a well-demarcated doublet in both reducing and non-reducing conditions in SDS-PAGE (Figure 4, lanes a and b). Two well-separated bands at 30-31KD are seen in the presence of 50 mM DTT (lane a), which migrate at 27-28KD in the absence of reducing agent (lane b). The clear-cut decrease in electrophoretic mobility after reduction with DTT strongly indicates that T11_{ex2} contains intrachain disulfide bridges; it does not form interchain bridges. Although not shown, microsequencing analysis of ³⁵S-cysteine labeled peptides verifies that there are two sets of intrachain disulfide bonds in T11_{ex2} between the amino-terminal cysteines and carboxy-terminal cysteines.

To investigate the difference between the two bands representing T11_{ex2}, 160 pmole of purified protein was separated by SDS-PAGE and blotted onto a PVDF membrane (Matsudaira, J. Biol. Chem. 262:10035-10038 (1987)). The upper and lower bands were cut separately from the membrane for amino-terminal sequencing. Each band yielded the CD2 amino-terminal sequence, suggesting that they differ from one another by post-translational modification.

-23-

As shown in Figure 4, endoglycosidase digestion generates at least five distinct bands. After short digestion times, two new lower molecular weight species are generated (lanes f-h). Some glycans on T11_{ex2} are apparently quite susceptible to digestion since even after simultaneous addition of enzyme and SDS sample buffer, these new species are generated (lane f). After 1 h digestion, most of the T11_{ex2} protein is digested to a 25KD species (lane i); overnight digestion results in complete digestion to a single band at 25KD (lane j). Note that the band of approximately 45KD size represents the Endo-F enzyme since it appears in lane k, where an equivalent amount of enzyme alone has been analyzed. The origin of the faint band at 48KD in lane j is unclear.

To determine whether T11_{ex2} exists as a non-covalently linked multimer in aqueous solution, it was subjected to equilibrium sedimentation by the high-speed meniscus depletion method (Yphantis, Biochem. 3:297-317 (1964) in both aqueous and dissociating conditions. As shown in Figure 5, the calculated molecular weights for both conditions are very similar (25.3KD in aqueous solution vs. 24.7KD in dissociating conditions). This result demonstrates that T11_{ex2} exists as a monomer in solution.

Figure 5 shows the equilibrium sedimentation analysis as a plot of log (fringe displacement) against square of distance from center of rotation, r^2 . T11_{ex2} (0.05%) was analyzed by sedimentation

equilibrium on a Beckman model E analytical ultracentrifuge in aqueous solution (PBS) at 30,000 rpm ($\omega = 3.142 \times 10^3$ rad/sec) or in dissociating conditions (6 M guanidine hydrochloride) at 44,000 rpm ($\omega = 4.608 \times 10^3$ rad/sec). Data were obtained at 22°C (PBS) or 23°C (guanidine-hydrochloride). Assuming a partial specific volume of 0.725, the calculated molecular weight from the displayed slope and using the equation: $MW = 4.606 RT \times \text{slope} / [\omega(1 - v^{\text{bar}} \omega)]$ is 25,315 daltons in aqueous solution. Assuming a partial specific volume (denoted as v^{bar}) of 0.725 - 0.1 in 6 M guanidine hydrochloride (Richardson, N.E., *et al.*, Biochem J. **135**:87-92 (1973)), the calculated molecular weight is 24,736 daltons in dissociating conditions.

The expression of CD2 epitopes was investigated by immunoprecipitation analysis. The T11_{ex2} molecule can be immunoprecipitated by both anti-T11₁ and a second monoclonal antibody to a different epitope termed anti-T11₂. However, T11_{ex2} is not immunoprecipitated by the anti-CD2 antibody, anti-T11₃, which defines an activation specific epitope on CD2. Nevertheless, T11_{ex2} was able to inhibit the binding of anti-T11₃-FITC to the T11₃ + Jurkat cell line at a concentration of 10 μ M, implying its presence on T11_{ex2} (data not shown). These results also suggest that the affinity of anti-T11₃ for its epitope is low.

Secondary structure predictions suggest the presence of both α -helical and β -sheet structure in the CD2 external domain (Clayton *et al.*, Eur. J. Immunol. **17**:1367-1370 (1987)). To more directly

-25-

predict secondary structural characteristics, the T11_{ex2} molecule was evaluated by circular dichroism. Figure 6 shows the circular dichroism spectra of T11_{ex2}. Far ultraviolet circular dichroism spectra represent the average of 3-5 individual spectra with data taken at 0.5 nm wavelength intervals in 10mM sodium phosphate, pH 7.2 A: spectrum at 25°C of untreated T11_{ex2}; B: spectrum of T11_{ex2} reduced with 10 μ M DTT and alkylated with 20 μ M iodoacetamide; C: thermal denaturation of the sample in A; D: thermal denaturation of the sample in B.

As shown in Figure 6A, the far ultraviolet CD spectrum of T11_{ex2} in 10 mM sodium phosphate shows a positive absorption maximum at about 200 nm ($\Delta\epsilon = 0.459$), a negative minimum at 215 nm ($\Delta\epsilon = -1.94$) and shoulder at 225 nm ($\Delta\epsilon = -1.0$). When the T11_{ex2} molecule is reduced by 50 mM DTT and subsequently alkylated with iodoacetamide, the CD spectrum is substantially altered, pointing to a role for disulfide bridges in stabilizing secondary and tertiary structure (Figure 6B). The fact that the spectrum of the non-reduced molecule reflects significant thermal denaturation at 80°C (Figure 6C) confirms that substantial secondary structure is present in soluble T11_{ex2}. As expected, the pattern after thermal denaturation is the same for the reduced as for the non-reduced molecule (compare Figures 6C and 6D).

In its overall pattern, the CD spectrum resembles that for Thy-1 (Campbell *et al.*, Nature 282:341-342 (1979)) which is a well-recognized

-26-

member of the immunoglobulin superfamily (Williams and Barclay, Annu. Rev. Immunol. 6:381-405 (1988)) and is therefore predicted to consist entirely of β -sheet. However, the shoulder at 225 nm is absent from the Thy-1 profile. To obtain a more objective prediction of secondary structure from the CD spectrum, the digitalized absorption data (Table I) were deconvoluted according to the inverse matrix method of Compton and Johnson, Anal. Biochem. 155:155-167 (1986). The resulting predictions for proportions of secondary elements are: α -helix, 20%; anti-parallel β -sheet, 13%; parallel β -sheet, 9%, turn, 20%, other, 46%. Since none of the protein in the data sets used to determine the matrix values are homologous to CD2, the predicted fractions of secondary structure are only approximate. When the same CD data are deconvoluted by the method of Yang et al., Meth. Enzymol. 130:208-209 (1986a), 15% α and 40% β structures are predicted.

-27-

TABLE I
 DIGITIZED PROTEIN CIRCULAR DICHROISM SPECTRUM
 FOR T11_{ex2} FROM 184-260 nm at 2 nm INTERVALS

<u>Wavelength</u>	<u>$\Delta\epsilon$</u>	<u>Wavelength</u>	<u>$\Delta\epsilon$</u>
184	-1.49	224	-1.002
186	-1.50	226	-0.923
188	-1.345	228	-0.856
190	-1.176	230	-0.743
192	-1.026	232	-0.629
194	-0.284	234	-0.515
196	0.241	236	-0.426
198	0.411	238	-0.348
200	0.459	240	-0.282
202	0.360	242	-0.234
204	0.027	244	-0.195
206	-0.432	246	-0.151
208	-0.959	248	-0.112
210	-1.467	250	-0.089
212	-1.750	252	-0.077
214	-1.949	254	-0.062
216	-1.930	256	-0.044
218	-1.771	258	-0.018
220	-1.515	260	-0.018
222	-1.241		

Spectrum was taken in 10 mM sodium phosphate (pH 7.2) in a 1 mm cell at 22°C. Secondary structure predictions were calculated by taking dot products with inverse CD spectra for the five secondary

-28-

structure categories as described in Table 6 of Compton and Johnson, Appl. Biochem. 155:155-167 (1986). The calculated results are: α -helix, 20%; anti-parallel β -sheet, 13%; parallel β -sheet, 9%; turn, 20%; other, 46%. Units for $\Delta\epsilon$ are $M^{-1}cm^{-1}$.

To investigate the presence of protease-resistant domains in the T11_{ex2} molecule, limited papain digestions were performed. Fifteen min. digestion at an enzyme:protein ratio of 1:250 at 37°C yields a major band at 15KD when analyzed by SDS-PAGE (data not shown). Longer digestion times cause partial disappearance of the 15KD band. Approximately 400 pmol of this 15KD material was blotted onto PVDF and sequenced, yielding 19 unambiguous residues corresponding to the CD2 amino-terminus. These results demonstrate that the carboxy region of T11_{ex2} is much less resistant to papain digestion than its amino-terminal counterpart.

T11_{ex2} inhibits sheep erythrocyte rosetting

The ability of T11_{ex2} to interact with the CD2 ligand expressed on the surface of various cell types was investigated. The capacity of T11_{ex2} to inhibit sheep erythrocyte rosetting with T lymphocytes was tested. Table II shows that rosetting is completely inhibited at concentrations of T11_{ex2} greater than 5 μM ; half-maximal inhibition occurs between 0.63 and 1.25 μM T11_{ex2}. Note that the anti-T11₁ (3T4-8B5) antibody abrogates rosetting at a concentration as low as 0.007 μM (Table II). This result suggests that any direct interaction between the soluble T11_{ex2} molecule and the CD2 ligand is of relatively low affinity.

-29-

TABLE II
CONCENTRATION DEPENDENCE OF T11_{ex2} INHIBITION OF
SHEEP ERYTHROCYTE-HUMAN T CELL ROSETTES

<u>Protein Added (μm)</u>		<u>Rosettes (% of Control)</u>
None		100
T11 _{ex2}	20	0
	10	0
	5	0
	2.5	4.8
	1.25	9.9
	0.63	122
	0.31	84
	0.16	116
	0.08	110
T4 _{ex1}	20	66
	10	91
	5	96
	2.5	105
	1.25	90
	0.63	72
	0.31	110
	0.16	99
	0.08	116
Anti-CD2	0.7	0
	0.07	0
	0.006	0

For rosette formation, sheep erythrocytes were preincubated with soluble protein or antibody at the indicated concentrations for 1 h at 4°C followed by the addition of Jurkat cells, co-centrifugation and a further 1 h incubation at 4°C. Rosettes were counted in a hemacytometer. Control value was the fraction of rosette formation in the absence of added protein. Anti-CD2 was the anti-T11₁ antibody of 3T4-8B5.

T11_{ex2} blocks the binding of anti-LFA-3 monoclonal antibody

To investigate whether T11_{ex2} could interact with a previously defined ligand for CD2 on human cells (Selveraj *et al.*, *Nature* 326:400-403 (1987), its ability to block the binding of monoclonal anti-LFA-3 antibody TS9/2 (Sanchez-Madrid *et al.*, *Proc. Natl. Acad. Sci. USA* 79:7489-7493 (1982) to LFA-3 bearing cells was tested. Specifically, anti-LFA-3 reactivity was measured by FACS analysis on the human B lymphoblastoid line JY, which expresses high levels of LFA-3. As shown in Table III, preincubation of JY cells with soluble CD4 does not affect this staining, whereas preincubation with 10 μ M T11_{ex2} causes a substantial decrease in observed fluorescence, reducing linear immunofluorescence from channel 120 to 30.6. This represents a reduction of 85% of specific anti-LFA-3 reactivity (calculated after subtraction of the background fluorescence of 13.2 linear units). Significant blocking is also seen at 1 μ M but not at 0.1 μ M. Addition of the control soluble CD4 T4_{ex1} protein has no effect on anti-LFA-3 binding. Thus, by this measure, as well as by inhibition of sheep erythrocyte rosetting, the affinity of T11_{ex2} for its ligand is apparently in the micromolar range.

- 31 -

TABLE III

CONCENTRATION DEPENDENCE OF T11_{ex2} INHIBITION OF
MONOCLONAL ANTI-LFA-3 ANTIBODY BINDING

<u>Protein Added</u>	<u>Anti-LFA-3 Antibody Added</u>	<u>Mean Fluorescence Intensity</u>
T11 _{ex2} :		
300 µg/ml (10 µM)	+	30.6
30 µg/ml (1 µM)	+	66.1
3 µg/ml (0.1 µM)	+	109.1
0.3 µg/ml (0.01 µM)	+	119.8
T4 _{ex1} :		
300 µg/ml	+	125.1
30 µg/ml	+	120.2
3 µg/ml	+	117.6
0.3 µg/ml	+	116.9
None	+	123.7
	-	13.2

Mean fluorescence intensities were obtained on a linear scale from 0-250.

T11_{ex2} binds to LFA-3 on human β lymphoblastoid cells

Since T11_{ex2} clearly inhibits sheep erythrocyte rosetting and blocks the binding of anti-LFA-3 antibody, we determined whether specific, saturable binding of T11_{ex2} to human cells bearing LFA-3 could be detected. Two types of binding assays were employed. In the first, increasing amounts of unlabeled T11_{ex2} was added to a mixture of the JY B lymphoblastoid cell line plus a constant amount of ¹²⁵I labeled T11_{ex2} (ligand).

Figure 7 shows the competitive inhibition of radioiodinated T11_{ex2} binding to JY cells. 5×10^6 cpm radiolabelled T11_{ex2} (2.8×10^8 cpm/nmole) was added at $0.1 \mu\text{M}$ to 1.8×10^6 JY cells overlayed onto 0.2 ml of a 1.5:1 mixture of dibutyl phthalate: dioctyl phthalate: dioctyl phthalate in 0.5 ml plastic tubes. Increasing concentrations of unlabeled T11_{ex2} (closed circles) or T4_{ex1} as a control (open circles) were added in a final volume of 200 μl in RPMI 1640/10% FCS. After 1 h incubation at 4°C, tubes were spun, cell pellets severed and bound radioactivity was determined. To one tube anti-LFA-3 antibody TS2/9 was added at 50 $\mu\text{g/ml}$ as a determination of non-specific binding. Specific cpm are the total cpm minus cpm bound in the presence of anti-LFA-3 (9070 cpm). Binding in the presence on control anti-HLA monoclonal antibody W6/32 is also shown (triangle).

As shown in Figure 7, binding of radiolabeled T11_{ex2} is progressively inhibited by the addition of

-33-

increasing amounts of unlabeled molecules.

Half-maximal inhibition occurs at about $0.5 \mu\text{M}$

Tl1_{ex2} . In contrast and as expected, addition of soluble CD4 T4_{ex1} protein has no effect on ^{125}I -labeled Tl1_{ex2} binding to JY.

As a second measure of binding, increasing amounts of labeled Tl1_{ex2} were added to a constant number of JY cells. Figure 8 shows the saturation binding of Tl1_{ex2} to JY cells and Scatchard analysis: Increasing concentrations of radiolabeled Tl1_{ex2} (1.31×10^7 cpm/nmole) were added to 2.6×10^6 JY cells in the presence or absence of $50 \mu\text{g/ml}$ anti-LFA-3 antibody to determine nonspecific binding. Binding was carried out as above and the dissociation constant determined by Scatchard analysis after subtraction of nonspecific binding determined in the presence of anti-LFA-3. Top: total binding (open circles) and specific binding (closed circles). Bottom: Scatchard analysis: correlation coefficient = 0.98; $-1/\text{slope} - K_d = 0.4 \mu\text{M}$.

As shown in Figure 8 (top), specific binding is saturable. These saturation binding data were transformed by Scatchard analysis (Figure 8, bottom). Specific Tl1_{ex2} binding is saturated at about 300×10^3 molecules per cell. The Scatchard plot yields an estimated dissociation constant of $0.4 \mu\text{M}$ for the Tl1_{ex2} -LFA-3 interaction. Note that although a single incubation period (1 h at 4°C) of cells with ligand was used in these experiments, kinetic analysis demonstrates that maximal Tl1_{ex2} -LFA-3 binding occurs within 5 min.

-34-

Conclusion

We have shown that a soluble, monomeric extracellular segment CD2 molecule, termed T11_{ex2}, carries epitopes recognized by antibodies against the native surface CD2 structure on resting T lymphocytes and interacts specifically with the surface-bound CD2 ligand, LFA-3. The measured dissociation constant for this interaction is 0.4 μ M, implying a low affinity relative to hormone receptor-ligand interactions (i.e., for IL-2 $K_d = 10^{-11}$ M) (Smith, 1984) but equivalent to that of primary antibody responses ($K_d = 10^{-5}$ to 10^{-6}) (Eisen and Siskind, 1964). The T11_{ex2} molecule gives rise to a proteolytically-resistant 15KD amino-terminal papain fragment, suggesting that the amino-terminal ~100 amino acid residues comprise a stable, well-folded domain corresponding to a polypeptide encoded by the first extracellular exon. In contrast, despite disulfide linkages, the carboxy-terminal encoded extracellular domain is labile to papain.

Deposit

The virus, AcNP/T11_{ex2} has been deposited at the American Type Culture Collection in Rockville, Maryland on January 19, 1989, and assigned the ATCC Accession Number _____.

-35-

Equivalents

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiment of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

- 36 -

CLAIMS

1. A soluble peptide having a lymphocyte function-associated antigen 3 (LFA-3) binding domain and antigenic epitopes recognized by antibodies raised against native CD2 on the surface of human T cells; the peptide being capable of binding LFA-3 and forming at least two intramolecular disulfide bonds.
2. A soluble peptide of Claim 1 wherein the antigenic epitopes are T11₁, T11₂ and T11₃.
3. A soluble peptide of Claim 2 wherein the peptide exists as a monomer in aqueous medium.
4. An isolated soluble peptide having an amino acid sequence encoded by the two extracellular segment exons of the gene encoding human CD2 glycoprotein, the peptide being capable of binding LFA-3, inhibiting CD2-mediated T cell activation and reacting with antibodies raised against native CD2 on the surface of human T cells.
5. A soluble peptide of Claim 4 wherein the peptide exists as a monomer in aqueous medium.

-37-

6. The soluble peptide of Claim 4 having the amino acid sequence:

Lys Glu Ile Thr Asn Ala Leu Glu Thr Trp
 Gly Ala Leu Gly Gln Asp Ile Asn Leu Asp
 Ile Pro Ser Phe Gln Met Ser Asp Asp Ile
 Asp Asp Ile Lys Trp Glu Lys Thr Ser Asp
 Lys Lys Lys Ile Ala Gln Phe Arg Lys Glu
 Lys Glu Thr Phe Lys Glu Lys Asp Thr Tyr
 Lys Leu Phe Lys Asn Gly Thr Leu Lys Ile
 Lys His Leu Lys Thr Asp Asp Gln Asp Ile
 Tyr Lys Val Ser Ile Tyr Asp Thr Lys Gly
 Lys Asn Val Leu Glu Lys Ile Phe Asp Leu
 Lys Ile Gln Glu Arg Val Ser Lys Pro Lys
 Ile Ser Trp Thr Cys Ile Asn Thr Thr Leu
 Thr Cys Glu Val Met Asn Gly Thr Asp Pro
 Glu Leu Asn Leu Tyr Gln Asp Gly Lys His
 Leu Lys Leu Ser Gln Arg Val Ile Thr His
 Lys Trp Thr Thr Ser Leu Ser Ala Lys Phe
 Lys Cys Thr Ala Gly Asn Lys Val Ser Lys
 Glu Ser Ser Val Glu Pro Val Ser Cys Pro
 Glu Lys

and modifications of the peptide in which amino acid residues have been deleted, inserted or substituted without essentially detracting from the properties thereof.

7. The soluble peptide of Claim 6 comprising T11₁, T11₂ and T11₃ epitopes localized on the peptide.

-38-

8. The soluble peptide of Claim 7 having the LFA-3 binding domain.
9. An isolated DNA sequence encoding a soluble peptide having a lymphocyte function-associated antigen 3 binding domain and antigenic epitopes recognized by antibodies raised against native CD2 on the surface of human T cells; the peptide being capable of binding LFA-3 and forming at least two intramolecular disulfide bonds.
10. An isolated DNA sequence encoding a soluble peptide having an amino acid sequence encoded by the two extracellular segment exons of the gene encoding human CD2 glycoprotein, the peptide being capable of binding LFA-3, inhibiting CD2-mediated T cell activation and reacting with antibodies raised against native CD2 on the surface of human T cells.
11. An isolated DNA sequence encoding the amino acid sequence of Claim 6 or substantial coding equivalents thereof.
12. An expression vector having an isolated DNA sequence as claimed in Claim 9.
13. An expression vector having the isolated DNA sequence as claimed in Claim 10.

-39-

14. An expression vector having an isolated DNA sequence encoding the amino acid sequence of Claim 5.
15. An expression vector of Claim 9 wherein the vector is a baculovirus transfer vector.
16. A cell transformed with the expression vector of Claim 12.
17. A cell transformed with the expression vector of Claim 13.
18. A cell transformed with the expression vector of Claim 14.
19. A plasmid having the ATCC Accession Number
-----.
20. A method of inhibiting T cell activation comprising the step of administering to a patient a soluble peptide having a lymphocyte function-associated antigen 3 binding domain and antigenic epitopes recognized by antibodies raised against native CD2 on the surface of human T cells; the peptide being capable of binding LFA-3 and forming at least two intramolecular disulfide bonds.

21. A method of Claim 20 wherein the soluble peptide is administered intravenously.
22. A method of inhibiting T cell activation comprising the step of administering to a patient a soluble peptide having an amino acid sequence encoded by the two extracellular segment exons of the gene encoding human CD2 glycoprotein, the peptide being capable of binding LFA-3, inhibiting T cell activation and reacting with antibodies raised against the native CD2 protein on the surface of human T cells.

23. A method of Claim 22 wherein the soluble peptide comprise the amino acid sequence:

Lys Glu Ile Thr Asn Ala Leu Glu Thr Trp
 Gly Ala Leu Gly Gln Asp Ile Asn Leu Asp
 Ile Pro Ser Phe Gln Met Ser Asp Asp Ile
 Asp Asp Ile Lys Trp Glu Lys Thr Ser Asp
 Lys Lys Lys Ile Ala Gln Phe Arg Lys Glu
 Lys Glu Thr Phe Lys Glu Lys Asp Thr Tyr
 Lys Leu Phe Lys Asn Gly Thr Leu Lys Ile
 Lys His Leu Lys Thr Asp Asp Gln Asp Ile
 Tyr Lys Val Ser Ile Tyr Asp Thr Lys Gly
 Lys Asn Val Leu Glu Lys Ile Phe Asp Leu
 Lys Ile Gln Glu Arg Val Ser Lys Pro Lys
 Ile Ser Trp Thr Cys Ile Asn Thr Thr Leu
 Thr Cys Glu Val Met Asn Gly Thr Asp Pro
 Glu Leu Asn Leu Tyr Gln Asp Gly Lys His
 Leu Lys Leu Ser Gln Arg Val Ile Thr His
 Lys Trp Thr Thr Ser Leu Ser Ala Lys Phe
 Lys Cys Thr Ala Gly Asn Lys Val Ser Lys
 Glu Ser Ser Val Glu Pro Val Ser Cys Pro
 Glu Lys

and modifications of the peptide in which amino acid residues have been deleted, inserted or substituted without essentially detracting from the properties thereof.

24. A method of Claim 23 wherein the soluble peptide is administered intravenously.

-24	Met	Ser	Phe	Pro	Cys	Lys	Phe	Val	Ala	Ser	Phe	Leu	Leu	Ile	Phe	L
-9	Asn	Val	Ser	Ser	Lys	G										
7	Leu	Glu	Thr	Trp	Gly	Ala	Ala	Val	Ser	Lys	Glu	Ile	Thr	Asn	Ala	Ex1
22	Pro	Ser	Phe	Gln	Met	Ser	Leu	Gly	Gln	Asp	Ile	Asn	Leu	Asp	Ile	
37	Lys	Thr	Ser	Asp	Lys	Lys	Lys	Asp	Ile	Gln	Asp	Ile	Lys	Trp	Glu	
52	Glu	Thr	Phe	Lys	Glu	Lys	Asp	Ile	Ala	Lys	Phe	Arg	Lys	Glu	Lys	
67	Thr	Leu	Lys	Ile	Lys	His	Leu	Thr	Tyr	Lys	Leu	Phe	Lys	Asn	Gly	
82	Lys	Val	Ser	Ile	Tyr	Asp	Thr	Lys	Thr	Asp	Asp	Gln	Asp	Ile	Tyr	
97	Ile	Phe	Asp	Leu	Lys	Ile	Gln	Lys	Gly	Lys	Asn	Val	Leu	Glu	Lys	
112	Ser	Trp	Thr	Cys	Ile	Asn	Thr	Leu	Arg	Val	Ser	Lys	Pro	Lys	Ile	Ex2
127	Gly	Thr	Asp	Pro	Glu	Leu	Asn	Leu	Leu	Thr	Cys	Glu	Val	Met	Asn	
142	Lys	Leu	Ser	Gln	Arg	Val	Ile	Thr	Tyr	Gln	Asp	Gly	Lys	His	Leu	
157	Ser	Ala	Lys	Phe	Lys	Cys	Thr	Ala	His	Lys	Trp	Thr	Thr	Ser	Leu	
172	Ser	Ser	Val	Glu	Pro	Val	Ser	Cys	Pro	Asn	Lys	Val	Ser	Lys	Glu	
187	Tyr	Leu	Ile	Ile	Gly	Ile	Cys	Gly	Val	Lys	Lys	Gly	Leu	Asp	Ile	TM
202	Phe	Val	Ala	Leu	Leu	Val	Phe	Tyr	Gly	Gly	Ser	Leu	Leu	Met	Val	
217	Arg	Ser	Arg	Arg	Asn	A			Ile	Thr	Lys	Arg	Lys	Lys	Gln	
232	Val	Ala	Thr	Glu	Glu	Arg	Glu	Gly	Leu	Glu	Thr	Arg	Ala	His	Arg	C
247	Ser	Thr	Pro	Gln	Asn	Pro	Gly	Arg	Lys	Pro	His	Gln	Ile	Pro	Ala	
262	Pro	Gly	His	Arg	Ser	Gln	Ala	Thr	Ser	Gln	His	Pro	Pro	Pro	Pro	
277	Gly	His	Arg	Val	Gln	His	Gln	Pro	Ser	His	Arg	Pro	Pro	Pro	Pro	
292	Ser	Gly	Thr	Gln	Val	His	Gln	Gln	Lys	Lys	Arg	Pro	Pro	Ala	Arg	
307	Pro	Arg	Val	Gln	Pro	Lys	Pro	Pro	Gly	Gly	Pro	Ala	Leu	Pro	Asn	
322	Leu	Ser	Pro	Ser	Ser	Asn	Pro	Pro	His	Gly	Ala	Glu	Glu	Asn	Ser	

FIG. 1

SUBSTITUTE SHEET

										38											53																																
										ATG AGC TTT CCA TGT AAA TTT GTA GCC AGC TTC																																											
										MET Ser Phe Pro Cys Lys Phe Val Ala Ser Phe																																											
										68											83											98																					
CTT CTG ATT TTC AAT GTT TCT TCC AAA GGT GCA GTC TCC AAA GAG ATT ACG AAT																																																					
Leu Leu Ile Phe Asn Val Ser Ser Lys Gly Ala Val Ser Lys Glu Ile Thr Asn																																																					
										113											128											143											158										
GCC TTG GAA ACC TGG GGT GCC TTG GGT CAG GAC ATC AAC TTG GAC ATT CCT AGT																																																					
Ala Leu Glu Thr Trp Gly Ala Leu Gly Gln Asp Ile Asn Leu Asp Ile Pro Ser																																																					
										173											188											203											218										
TTT CAA ATG AGT GAT GAT ATT GAC GAT ATA AAA TGG GAA AAA ACT TCA GAC AAG																																																					
Phe Gln MET Ser Asp Asp Ile Asp Asp Ile Lys Trp Glu Lys Thr Ser Asp Lys																																																					
										233											248											263																					
AAA AAG ATT GCA CAA TTC AGA AAA GAG AAA GAG ACT TTC AAG GAA AAA GAT ACA																																																					
Lys Lys Ile Ala Gln Phe Arg Lys Glu Lys Glu Thr Phe Lys Glu Lys Asp Thr																																																					
										278											293											308											323										
TAT AAG CTA TTT AAA AAT GGA ACT CTG AAA ATT AAG CAT CTG AAG ACC GAT GAT																																																					
Tyr Lys Leu Phe Lys Asn Gly Thr Leu Lys Ile Lys His Leu Lys Thr Asp Asp																																																					
										338											353											368																					
CAG GAT ATC TAT AAG GTA TCA ATA TAT GAT ACA AAA GGA AAA AAT GTG TTG GAA																																																					
Gln Asp Ile Tyr Lys Val Ser Ile Tyr Asp Thr Lys Gly Lys Asn Val Leu Glu																																																					
										383											398											413											428										
AAA ATA TTT GAT TTG AAG ATT CAA GAG AGG GTC TCA AAA CCA AAG ATC TCC TGG																																																					
Lys Ile Phe Asp Leu Lys Ile Gln Glu Arg Val Ser Lys Pro Lys Ile Ser Trp																																																					
										443											458											473											488										
ACT TGT ATC AAC ACA ACC CTG ACC TGT GAG GTA ATG AAT GGA ACT GAC CCC GAA																																																					
Thr Cys Ile Asn Thr Thr Leu Thr Cys Glu Val MET Asn Gly Thr Asp Pro Glu																																																					
										503											518											533																					
TTA AAC CTG TAT CAA GAT GGG AAA CAT CTA AAA CTT TCT CAG AGG GTC ATC ACA																																																					
Leu Asn Leu Tyr Gln Asp Gly Lys His Leu Lys Leu Ser Gln Arg Val Ile Thr																																																					
										548											563											578											593										
CAC AAG TGG ACC ACC AGC CTG AGT GCA AAA TTC AAG TGC ACA GCA GGG AAC AAA																																																					
His Lys Trp Thr Thr Ser Leu Ser Ala Lys Phe Lys Cys Thr Ala Gly Asn Lys																																																					
										608											623											638																					
GTC AGC AAG GAA TCC AGT GTC GAG CCT GTC AGC TGT CCA GAG AAA GGT CTG GAC																																																					
Val Ser Lys Glu Ser Ser Val Glu Pro Val Ser Cys Pro Glu Lys Gly Leu Asp																																																					
										653											668											683											698										
ATC TAT CTC ATC ATT GGC ATA TGT GGA GGA GGC AGC CTC TTG ATG GTC TTT GTG																																																					
Ile Tyr Leu Ile Ile Gly Ile Cys Gly Gly Gly Ser Leu Leu MET Val Phe Val																																																					
										713											728											743											758										
GCA CTG CTC GTT TTC TAT ATC ACC AAA AGG AAA AAA CAG AGG AGT CGG AGA AAT																																																					
Ala Leu Leu Val Phe Tyr Ile Thr Lys Arg Lys Lys Gln Arg Ser Arg Arg Asn																																																					
										773											788											803																					
GAT GAG GAG CTG GAG ACA AGA GCC CAC AGA GTA GCT ACT GAA GAA AGG GGC CGG																																																					
Asp Glu Glu Leu Glu Thr Arg Ala His Arg Val Ala Thr Glu Glu Arg Gly Arg																																																					

FIG. 1A

SUBSTITUTE SHEET

818 833 848 863
 AAG CCC CAC CAA ATT CCA GCT TCA ACC CCT CAG AAT CCA GCA ACT TCC CAA CAT
 Lys Pro His Gln Ile Pro Ala Ser Thr Pro Gln Asn Pro Ala Thr Ser Gln His 257
 878 893 908
 CCT CCT CCA CCA CCT GGT CAT CGT TCC CAG GCA CCT AGT CAT CGT CCC CCG CCT
 Pro Pro Pro Pro Pro Gly His Arg Ser Gln Ala Pro Ser His Arg Pro Pro Pro 275
 923 938 953 968
 CCT GGA CAC CGT GTT CAG CAC CAG CCT CAG AAG AGG CCT CCT GCT CCG TCG GGC
 Pro Gly His Arg Val Gln His Gln Pro Gln Lys Arg Pro Pro Ala Pro Ser Gly 293
 983 998 1013 1028
 ACA CAA GTT CAC CAG CAG AAA GGC CCG CCC CTC CCC AGA CCT CGA GTT CAG CCA
 Thr Gln Val His Gln Gln Lys Gly Pro Pro Leu Pro Arg Pro Arg Val Gln Pro 311
 1043 1058 1073
 AAA CCT CCC CAT GGG GCA GCA GAA AAC TCA TTG TCC CCT TCC TCT AAT
 Lys Pro Pro His Gly Ala Ala Glu Asn Ser Leu Ser Pro Ser Ser Asn 327

FIG. 1A (CONT)

SUBSTITUTE SHEET




FIG. 2




FIG. 3

SUBSTITUTE SHEET




FIG. 4




FIG. 5




FIG. 6




FIG. 7

SUBSTITUTE SHEET








FIG. 8

SUBSTITUTE SHEET

INTERNATIONAL SEARCH REPORT

International Application No PCT/US 89/00218

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) * According to International Patent Classification (IPC) or to both National Classification and IPC IPC ⁴ : C 12 N 15/00, C 12 P 21/04, C 07 K 13/00											
II. FIELDS SEARCHED <div style="display: flex; justify-content: space-between; margin-top: 10px;"> <div style="width: 40%;"> Classification System : IPC⁴ </div> <div style="width: 50%;"> Minimum Documentation Searched : Classification Symbols C 07 K, C 12 N, C 12 P </div> </div> <div style="margin-top: 10px;"> Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched : </div>											
III. DOCUMENTS CONSIDERED TO BE RELEVANT* <table border="1" style="width: 100%; border-collapse: collapse; margin-top: 5px;"> <thead> <tr> <th style="width: 10%;">Category *</th> <th style="width: 70%;">Citation of Document, ** with indication, where appropriate, of the relevant passages **</th> <th style="width: 20%;">Relevant to Claim No. **</th> </tr> </thead> <tbody> <tr> <td style="text-align: center; vertical-align: top;">A</td> <td>Chemical Abstracts, vol. 106, no. 21, 25 May 1987 (Columbus, Ohio, US) P. Selvaraj et al.: "The T lymphocyte glycoprotein CD2 binds the cell surface ligand LFA-3", see page 549, abstract no. 174287w & Nature (London) 1987, 326(6111) 400-3</td> <td style="text-align: center; vertical-align: top;">1,4,9,10</td> </tr> <tr> <td style="text-align: center; vertical-align: top;">A</td> <td>Chemical Abstracts, vol. 107, no. 15, 12 October 1987 (Columbus, Ohio, US) P.H. Sayre et al.: "Molecular cloning and expression of T11 cDNAs reveal a receptor-like structure on human T lymphocytes", see page 177, abstract no. 128218x & Proc. Natl. Acad. Sci. U.S.A., 1987, 84(9) 2941-5</td> <td style="text-align: center; vertical-align: top;">1-19</td> </tr> </tbody> </table>			Category *	Citation of Document, ** with indication, where appropriate, of the relevant passages **	Relevant to Claim No. **	A	Chemical Abstracts, vol. 106, no. 21, 25 May 1987 (Columbus, Ohio, US) P. Selvaraj et al.: "The T lymphocyte glycoprotein CD2 binds the cell surface ligand LFA-3", see page 549, abstract no. 174287w & Nature (London) 1987, 326(6111) 400-3	1,4,9,10	A	Chemical Abstracts, vol. 107, no. 15, 12 October 1987 (Columbus, Ohio, US) P.H. Sayre et al.: "Molecular cloning and expression of T11 cDNAs reveal a receptor-like structure on human T lymphocytes", see page 177, abstract no. 128218x & Proc. Natl. Acad. Sci. U.S.A., 1987, 84(9) 2941-5	1-19
Category *	Citation of Document, ** with indication, where appropriate, of the relevant passages **	Relevant to Claim No. **									
A	Chemical Abstracts, vol. 106, no. 21, 25 May 1987 (Columbus, Ohio, US) P. Selvaraj et al.: "The T lymphocyte glycoprotein CD2 binds the cell surface ligand LFA-3", see page 549, abstract no. 174287w & Nature (London) 1987, 326(6111) 400-3	1,4,9,10									
A	Chemical Abstracts, vol. 107, no. 15, 12 October 1987 (Columbus, Ohio, US) P.H. Sayre et al.: "Molecular cloning and expression of T11 cDNAs reveal a receptor-like structure on human T lymphocytes", see page 177, abstract no. 128218x & Proc. Natl. Acad. Sci. U.S.A., 1987, 84(9) 2941-5	1-19									
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>* Special categories of cited documents: **</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 50%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"A" document member of the same patent family</p> </div> </div>											
IV. CERTIFICATION <table border="1" style="width: 100%; border-collapse: collapse; margin-top: 5px;"> <tr> <td style="width: 50%; padding: 5px;"> Date of the Actual Completion of the International Search <div style="text-align: center;">10th August 1989</div> </td> <td style="width: 50%; padding: 5px;"> Date of Mailing of this International Search Report <div style="text-align: center;">- 1 SEP 1989</div> </td> </tr> <tr> <td style="width: 50%; padding: 5px;"> International Searching Authority <div style="text-align: center;">EUROPEAN PATENT OFFICE</div> </td> <td style="width: 50%; padding: 5px;"> Signature of Authorized Officer <div style="text-align: center;">  P.C.G. VAN DER PUTTEN </div> </td> </tr> </table>			Date of the Actual Completion of the International Search <div style="text-align: center;">10th August 1989</div>	Date of Mailing of this International Search Report <div style="text-align: center;">- 1 SEP 1989</div>	International Searching Authority <div style="text-align: center;">EUROPEAN PATENT OFFICE</div>	Signature of Authorized Officer <div style="text-align: center;">  P.C.G. VAN DER PUTTEN </div>					
Date of the Actual Completion of the International Search <div style="text-align: center;">10th August 1989</div>	Date of Mailing of this International Search Report <div style="text-align: center;">- 1 SEP 1989</div>										
International Searching Authority <div style="text-align: center;">EUROPEAN PATENT OFFICE</div>	Signature of Authorized Officer <div style="text-align: center;">  P.C.G. VAN DER PUTTEN </div>										

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)

Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
A	Chemical Abstracts, vol. 108, no. 1, 4 January 1988 (Columbus, Ohio, US) A. Peterson et al.: "Monoclonal antibody and ligand binding sites of the T cell erythrocyte receptor (CD2)" see page 427, abstract no. 4386b & Nature (London) 1987, 329(6142), 842-6 --	1,4,7,8-10 -
T	Chemical Abstracts, vol. 110, no. 17, 24 April 1989 (Columbus, Ohio, US) P.H. Sayre et al.: "Structural and binding analysis of a two domain extracellular CD2 molecule" see page 549, abstract no. 152376e & Exp. Med. 1989, 169(3), 995-1009 -----	1-19

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

V. ☒ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☒ Claim numbers XX because they relate to subject matter not required to be searched by this Authority, namely:

xx Claims 20-22

See PCT Rule 39-1(iv):

Methods for treatment of the human or animal body, by means of surgery of therapy, as well as diagnostic methods.

2. ☐ Claim numbers because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claim numbers because they are dependent claims and are not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☐ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING

This International Searching Authority found multiple inventions in this international application as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.
2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:
3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:
4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.